

Lao People's Democratic Republic

Ministry of Health

Center for HIV/AIDS/STI (CHAS)



Integrated Behavioral Biological Surveillance 2009

Supported by: Global fund to fight AIDS/TB/Malaria



USAID
FROM THE AMERICAN PEOPLE



Family Health
International



Investing in our future

The Global Fund
To Fight AIDS, Tuberculosis and Malaria



UNAIDS
JOINT UNITED NATIONS PROGRAMME ON HIV/AIDS

UNHCR
UNICEF
WFP
UNDP
UNFPA

UNODC
ILO
UNESCO
WHO
WORLD BANK

In July 2011, FHI became FHI 360.



FHI 360 is a nonprofit human development organization dedicated to improving lives in lasting ways by advancing integrated, locally driven solutions. Our staff includes experts in health, education, nutrition, environment, economic development, civil society, gender, youth, research and technology – creating a unique mix of capabilities to address today's interrelated development challenges. FHI 360 serves more than 60 countries, all 50 U.S. states and all U.S. territories.

Visit us at www.fhi360.org.

3.6. Survey procedures:

→ *Informed consent*

Upon recruitment, potential participants were explained the objectives and procedures of the study including: the roles of each of the different teams members, and the procedures for maintaining anonymity. The study team supervisor and one interviewer then proceeded to a private location where they read the detailed consent form written in Lao to the participant, answered any questions raised, and obtained oral informed consent that was witnessed. Potential participants were administered witnessed oral informed consent instead of written consent to ensure that no name was recorded. Each participant was given a list with names and contact of survey managers, and copies of the consent form was also available for participants who wished to keep them for reference.

→ *Interviews*

Data were collected through face-to-face interview conducted by trained same sex interviewer in a private location. The questionnaire was administered in Lao language that is spoken by all the populations and in all the provinces. Interview forms were cross-checked by the team supervisors in the field to ensure their completeness and were delivered to the provincial supervisor at the end of each day. They were kept in a secured location until they are delivered to the central supervisor in charge of securing them in the capital.

→ *Data entry*

Data were double entered into Epi-data, however, a single of those two entries was used for analysis.

3.7. Ethical review:

The National Ethics Committee for Health Research (NECHR) has given the Ethical Clearance, no 257/NECHR, dated 13 August 2009 for the study of “ Behavioral Surveillance Survey among Service Women in Five Provinces of Lao PDR.



IV. Results

Data were collected in August 2009 among 912 service women. All interviews were conducted in Lao language except for Attapeu where 18 interviews (12% of interviews in Attapeu province) were conducted in Vietnamese. Cluster information were not completed and data weighting was therefore not performed in the subsequent analysis.

4.1. Demographic characteristics

While the majority participants from Vientiane capital and Attapeu were recruited in night-clubs, most participants from other provinces worked in drinking shops. Service women were young: about half were teenagers (46%) and their mean age was 21 years old. The majority (78%) had never married and another 19% were divorced or separated. They had a low level of education with 84% who had not studied further than primary school. In Vientiane about half of the service women were native from Vientiane municipality. In other provinces only a minority of the service women were native from the province where they worked. Luang Prabang province provided a substantial proportion of service women to Northern provinces. Women from Vientiane provided about a quarter of the service women to the Southern provinces. Service women are mobile as 70% had been living in the province of interview for less than 6 months.

Table 2: Demographic characteristics of service women, by province

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Place of recruitment						
Night club	44	11	1	9	53	27
Drinking shop	46	77	64	85	21	57
Restaurant	7	2	32	0	11	10
Guest house/hotel	2	10	3	0	3	3
Karaoke	2	0	0	6	13	4
Age						
15-19	44	67	63	34	25	46
20-24	45	26	29	38	36	36
>=25	11	7	8	28	39	17
Mean	21	19	20	22	24	21
Median	20	19	19	21	23	20
Province of birth						
Vientiane	47	1	4	36	23	26
Luang Namtha	1	5	7	1		2
Bokeo	0	3	18	1		4
Saravane	0			4	1	1
Attapeu	0			1	10	2
Luang Prabang	17	44	30	13	6	21
Other	34	47	40	46	47	41
Thailand / Vietnam	1				13	2
Highest level of education attended						
<= Primary school	77	93	88	85	84	84
Lower secondary school	20	7	13	14	15	15
Upper secondary / university	3	0	0	1	1	1
Marital status						
Never married	83	84	82	70	65	78
Currently married	1	3	1	7	5	3
Divorced / widow	17	13	16	23	29	19
Language						
Lao	100	100	100	100	88	98
Vietnamese	0	0	0	0	12	2
Duration living in this city						
<=6 months	57	81	77	79	66	70
>6 months	43	19	23	21	34	30

4.2. First sex

Service women had first sex on average at 17 years of age. Depending on the province, a quarter to a third of the service women had initiated sex with a client.

Table 3: Age at first sex and first sexual partner among service women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Age at first sex						
Mean	17	16	16	18	18	17
Median	17	16	16	18	18	17
First sexual partner						
Husband	10	13	12	21	25	15
Boy friend	60	51	51	43	51	53
Client	24	31	30	33	22	27
Other	5	5	8	3	1	5

4.3. Reproductive history, contraceptive use and abortions

While 17% of service women currently had children, 10% had been pregnant in the past 6 months and another 10% were pregnant in the past 6 to 12 months. It is likely that most of the recent pregnancies were unwanted as 18% of the service women had undergone an abortion in the past year. The majority of the abortions were conducted by the women themselves, without any medical supervision. Overall, 15% of the service women reported using modern contraceptives other than condoms.

Table 4: Reproductive history, use of contraceptives and history of abortion among service women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Currently have children	12	13	13	23	29	17
Time since last pregnancy						
Never pregnant	64	63	72	58	52	62
<= 6 mth	16	10	5	10	5	10
>6 & <=12 months	6	12	11	12	11	10
>12 months	14	15	12	20	32	18
Ever aborted (among those ever pregnant)	79	75	67	64	68	71
Aborted past year (among all women)	21	18	16	16	16	18
Who conducted last abortion						
Self inflicted	67	83	64	34	55	61
Medical staff	33	17	36	61	45	38
Friend	0	0	0	5	0	1
Type of contraceptive currently used						
Pill	8	45	2	15	12	15
Injectable	0	11	1	5	3	3
IUD	0	0	0	3	1	1
Withdrawal	17	11	5	4	10	11
Currently using any modern contraceptive method appart from condom	8	45	3	16	12	15

4.4. History of selling sex

About a quarter of service women sold sex at sexual initiation. It is noteworthy that almost all (98%) reported having sold sex in the past 3 months, which qualifies the population as female sex workers. The turnover of service women was high, ranging from half of the service women being new to the job every 4 months in Bokeo, to half of the service women being new to the job every 12 months in Vientiane. Overall, a quarter of the service women had been doing this job for 24 months or more (data not shown). This high turnover means that most of the service women are new to comers in the sex trade market and are ready to get infected.

Depending on the province, half of the service women had been staying from 1 to 4 months in the same establishment. However, the mean was substantially higher than the median indicating that some women stay for a long period of time in the same establishment. Still this information indicates a high mobility of a fringe of

the service women population, which poses a challenge for outreach interventions and requires frequent visits to establishments to ensure a high coverage.

Table 5: History of selling sex and mobility of service women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Sold sex at first sex	24	31	30	33	22	27
Duration selling sex (months)						
Mean	15	15	12	12	18	14
Median	9	8	4	8	12	8
Sold sex in past 3 months	98	100	97	94	99	98
Had worked as service women in other provinces	8	34	25	33	40	25
Duration in current establishment (months)						
Mean	9	4	4	3	9	6
Median	4	1	1	1	3	2
Number of months doing this job						
Mean	15	15	12	12	18	14
Median	9	8	4	8	12	8

4.5. Sexual Partners

The mean number of clients reported by service women for the past week remains low and comparable to previous round of surveillance (mean: 2.5 clients per week). About half of service women reported having some regular partners paying with gifts, which are called “regular clients” in the present report. In addition, 14% had casual partners defined as non-paying and non-regular partners.

Table 6: Number and type of partners reported by service women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Had regular partner giving gift but no money in past 3 months	51	56	44	36	39	46
Number of clients past week						
Mean	2.3	3.4	2.4	2.4	2.3	2.5
Median	2	3	2	2	2	2
Had non-paying & non-regular partner past 3 months	18	13	11	16	7	14

4.6. Last clients characteristics

Clients of service women are a majority of Lao national (78%) or citizens from the neighboring Asian countries (21%). About half of service women did not know the profession of their last client. Civil servants represented 13% of the last clients, so did military and police. Sexual services are more expensive in Vientiane or Attapeu than in other cities. Half of the clients pay at least 250,000 Kips per sexual service.

Table 7: Characteristics of last client

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Nationality of last client						
Lao	71	77	84	89	76	78
Thai	7	2	7	2	1	4
China	9	20	7	5	0	8
Vietnam	1	1	1	4	22	5
Burma	9	0	0	0	0	3
Japan/Korea	2	0	0	0	0	1
Western	1	0	0	0	1	0
Profession of last client						
Gov staff	7	17	12	15	22	13
Military/Police	9	20	18	14	9	13
Salesman	9	9	7	11	14	10
Tourist	5	5	1	1	0	3
Driver	1	6	4	3	1	3
Company/factory	4	5	5	17	11	8
NGO staff	3	0	1	1	2	2
Don't know/ No answer	61	38	51	39	41	49
How much did last client pay (Kip)						
Mean	452,000	179,000	289,000	237,900	343,000	325,000
Median	300,000	150,000	250,000	200,000	300,000	250,000

4.7. Availability and use of male latex condoms

More than half of the service women had purchased condoms in the month prior interview. However less than 10% of service women were carrying condoms at the time of interview, except for Attapeu, where 44% of participants could show a condom to interviewers. In every surveyed province, the overwhelming majority reported that condoms were available at their workplace.

Table 8: Condom availability

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Carries and can show condom at time of interview	7	8	7	11	44	14
Purchased condom pasts month	48	78	68	80	51	62
Cost of a pack of 12 condoms (Kip)						
Mean	5,396	2,513	4,340	4,670	4,174	4,282
Median	5,000	2,000	5,000	3,000	5,000	5,000
Condom available at work place	90	83	93	79	76	85

In every province, more than 95% of service women reported using condom at last sex with client. Overall, 89% reported always using condoms with clients in the past week and 83% in the past month. However, they were 53% to report having clients refusing to use condoms in the past month. In most instances, clients were opposed to using condom: 39% of service women had clients who propose to pay more for unprotected sex, and 36% of service women had clients threatening them to obtain unprotected sex. Clients' difficulty to sustain an erection was a minor cause of unprotected sex: 12% of service women reported unprotected sex past month due to clients' alcohol imbibitions, whereas 2% of service women had unprotected sex in the past month caused by clients high on drugs.


While 81% had used condom at last sex with casual partner, less than half had used condom at last sex with regular client. Overall, 30% of service women had unprotected sex with a regular client in the past month and 4% with casual partners.

Table 9: Condom use

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Condom use at last sex						
With client	95	98	97	97	98	97
With regular client	50	54	49	48	42	49
With casual partner	71	90	76	91	90	81
Condom use with client past week						
Always	84	92	86	95	94	89
Sometime	12	8	9	4	6	9
Never	3	0	4	1	1	2
Condom use with client past month						
Always	79	85	77	89	91	83
Sometime	19	15	22	10	8	16
Never	2	0	1	1	1	1
Had client refusing to use condom past month	57	60	46	37	59	53
Had client proposing money for unprotected sex past month	43	53	34	27	36	39
Had been threatened by client for unprotected sex past month	38	43	19	24	55	36
Had client too drunk to use condom past month	11	24	8	7	8	12
Had client too high on drugs to use condom past month	1	3	1	3	1	2
Had client who removed condom before the end of intercourse past month	1	0	1	0	0	0
Condom use with casual partners past 3 months						
No casual partner	81	87	89	79	93	85
Always	12	12	8	18	6	11
Sometime	3	1	1	2	1	2
Never	4	1	3	1	0	2
Condom use with regular client past 3 months						
No regular client	49	44	56	66	61	54
Always	16	19	16	15	11	16
Sometime	18	19	14	6	11	14
Never	17	17	14	13	17	16
Had condom breakage past 3 months	26	30	14	28	23	24
Used two condoms on the top of one another at last sex	9	2	5	9	5	7
Used two condoms on the top of one another past 3 months	32	15	15	30	27	25

Service women were 24% to have experienced condom breakages in the past three month. They were 25% to have used two condoms on the top of one another in the past three months. Using two condoms at a time was associated with breakages as those who had used two condoms at a time in the past three months had experienced twice more breakages than those who had not used two condoms at a time in the past three months ($p < 0.001$, see Figure 1).

Figure 1: Condom breakage past 3 months by use of two condoms at a time



4.8. Female condom

While 76% of service women from Vientiane had heard of female condoms, only a quarter to a third of service women from province had heard of such device. In Vientiane, 20% of service women had ever used a female condom. In the provinces, however, less than 10% of service women had ever used a female condom. Among those who had ever used female condoms, 56% reported feeling comfortable using them.



Integrated Behavioral Biological Surveillance 2009



USAID
FROM THE AMERICAN PEOPLE



Investing in our future
The Global Fund
To Fight AIDS, Tuberculosis and Malaria



UNAIDS
JOINT UNITED NATIONS PROGRAMME ON HIV/AIDS

UNHCR
UNICEF
WFP
UNDP
UNFPA
UNODC
ILO
UNESCO
WHO
WORLD BANK

Table 10: Use of female condoms

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Ever heard of female condom	76	32	28	28	39	46
Ever used female condom	20	6	3	7	4	10
Ever used female condom (n)	n=59	n=8	n=5	n=11	n=6	n=89
Feel comfortable using female condom	56	50	60	73	33	56

4.9. Violence to service women

A quarter of the service women from Vientiane and a third of those in Luang Namtha had been forced into sex in the past year. In addition 5% of service women had been molested by clients in the past 3 months.

Table 11: Report of aggressions

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Forced into sex past year	25	32	16	11	15	21
Molested by client in the past 3 months	7	8	2	5	2	5

4.10. Drug use

Only 5% of service women had ever used opium, heroin or methamphetamine. One percent of service women acknowledge having injected drugs in the past year.

Table 12: Type of drug use among Service Women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Ever used opium, heroin or methamphetamine	4	7	8	4	7	5
Ever used methamphetamine	4	7	8	4	7	5
Ever used thinner	0	0	1	1	0	0
Ever used heroin	0	0	1	1	0	0
Ever used opium	0	0	0	1	0	0
Ever used cannabis	0	0	1	1	0	0
Injected drugs past year	1	0	1	2	0	1

4.11. Knowledge of HIV

The overwhelming majority of service women knew that HIV is transmitted through unprotected sex (98%), stained needles (95%), or from mother to child during pregnancy or delivery (92%). However, fewer reported that HIV could be avoided by being faithful to an uninfected faithful partner (78%) or by abstaining from sex (26%). It is plausible that service women reported what they thought as a possible strategy to protect themselves against HIV rather than reporting their knowledge of the strategy.

Misconceptions on HIV modes of transmission were still substantial: 36% believed that HIV could be transmitted through mosquito bites (52% in Saravanne which is the province most affected by malaria), 12 % believed they could identify HIV infected individuals, and another 12% believed that HIV could be transmitted by sharing food utensils. Moreover 38% believed that they could prevent from getting infected with HIV by taking antibiotics before or after sex.

Overall, 45% of service women had answered correctly to the five questions composing the UNGASS indicator (including: condom use, faithfulness, recognizing infected people, sharing food, and transmission from mosquitoes).

Table 13: Knowledge of HIV transmission and prevention among service women

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Know that HIV can be prevented by using condoms when having sex	99	100	97	96	98	98
Know that HIV can be prevented by being faithful to one partner	78	89	81	66	79	78
Know that HIV can be avoided by abstaining from sex	35	19	16	28	21	26
Know that HIV can be transmitted by stained needles	95	99	94	88	98	95
Know that HIV can be transmitted from mother to child during pregnancy	95	91	92	83	97	92
Know that HIV infected person can look healthy	87	96	81	82	93	88
Know that HIV is not transmitted by mosquitoes	71	63	66	48	68	64
Know that HIV is not transmitted by sharing food utensils with HIV infected person	90	87	84	82	95	88
Know that taking antibiotics before / after sex does not prevent HIV	73	67	80	59	83	72
UNGASS knowledge indicator	47	53	46	29	47	45

2.12. Reported symptoms of STI and attendance to STI clinic

Overall, 31% of service women reported having had STI symptoms in the past three months and 18% reported having symptoms at time of interview. The frequency of reporting of STI symptoms was higher in Vientiane and the Northern provinces than in the Southern provinces (Saravane and Attapeu). The most frequently reported symptom of STI was vaginal discharge that is known to be a poor predictor of STI. Genital ulcer, genital vesicles, and genital warts were seldom reported.


In Vientiane, about half of the service women had undergone a routine STI checkup in the past three months, and a third had attended the STI clinic twice or more. In the other surveillance provinces, less than 20% had a routine STI checkup in the past three months and 4% or less had attended multiple times.

Table 14: Reported symptoms of STI and attendance to routine checkups

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Currently have symptom of STI, including:	19	20	25	17	6	18
Lekorrhea	18	19	25	16	6	17
Genital ulcer	2	1	1	2	0	1
Genital vesicules	2	1	1	1	0	1
Genital wart	0	1	0	1	0	0
Had STI symptoms in the past 3 months, including:	35	40	36	23	17	31
Lekorrhea	34	39	36	21	16	30
Genital ulcer	5	1	1	2	1	3
Genital vesicules	5	1	1	1	0	2
Genital wart	1	1	0	1	0	1
Number of attendings to STI checkup in the past 3 months						
Never attended	52	88	86	90	81	75
One	15	8	10	7	17	12
2 to 3 times	22	3	3	3	1	9
>3 times	10	1	1	0	0	4

About of half of those from Vientiane who had symptoms of STI in the past three months sought treatment at a drop-in center. In other provinces the majority self treated of chose to ask for advises at a pharmacy.

Figure 2: Source of treatment at last episode of STI, which occurred in the past three months



4.13. Uptake of HIV testing services

The proportion of service women who acknowledged having ever been tested for HIV was in the range of 43% in Vientiane to 16% in Bokeo. While a third of service women from Vientiane and a quarter of the service women from the Southern provinces (Saravane and Attapeu) reported having received HIV test

Table 15: Uptake of HIV testing services

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Ever tested for HIV	43	23	16	34	35	33
Tested for HIV and received test results in the past 12 months	36	6	10	24	26	23

Service women were asked which type of services they received at last HIV test and their satisfaction with services provision. VCT services appear fairly comprehensive in all surveyed provinces but Luang Namtha. In Luang Namtha, the majority of those tested (80%) reported not receiving pre-test or post-test counseling and received not explanation about their test results.

Table 16: Array of services received by service women at last HIV testing

	Vientiane n=130 %	Luang Namtha n=35 %	Bokeo n=25 %	Saravane n=54 %	Attapeu n=53 %	Total n=297 %
Ever tested for HIV						
Had pre-test counseling at last test	98	20	88	89	98	86
Had post-test counseling at last test	96	20	76	85	100	84
Found the staff friendly	100	23	88	80	100	86
Received results the same day	99	23	88	83	100	87
Received information about HIV transmission	68	14	60	63	53	57
Received information on how to prevent HIV	95	17	84	93	98	85
Received explanation about the meaning of the test result	96	20	84	94	98	86
Received information on hygiene for people living with HIV	89	20	84	93	96	82

4.14. Source of information and coverage of prevention services

Overall, 77% of service women received information on HIV in the past year and there was no major difference across provinces. In Vientiane, 77% of service women acknowledge having received information on HIV from NGO in the past three months. In other provinces NGO coverage was substantially lower, ranging from 55% in Bokeo to 23% in Attapeu where there is no NGO implementing prevention program. In all survey provinces, the majority of service women had received free condoms in the past three months. However, only in Vientiane was the condom distribution taking place on a regular basis (more than half of service women had received free condoms on at least 3 occasions).

Most service women had received information on HIV and condoms when they started working at their current location. In most instances, the information was provided by colleagues (46%), the mamasan (22%), or outreach workers (21%).


Table 17: Coverage of prevention services

	Vientiane n=300 %	Luang Namtha n=150 %	Bokeo n=152 %	Saravane n=160 %	Attapeu n=150 %	Total n=912 %
Attended health staff meeting or outreach session on HIV past year	81	82	72	62	83	77
Received information on HIV from NGO in the past 3 months	74	38	55	39	23	51
Number of times received free condoms in past 3 months						
Never	18	21	36	45	25	28
Once	9	13	25	20	31	18
2 to 3 times	20	46	28	23	29	28
>3 times	53	19	11	13	15	27
Received printed HIV information material in the past 6 months	44	39	54	28	42	42
Have poster on HIV / STI at workplace	27	27	43	25	25	29
Received information on condom and HIV when initiated working at current place	74	96	59	61	83	74
Who provided information when initiating working at current workplace						
Colleague	36	72	44	31	44	46
Mamasan	18	17	16	36	27	22
Outreach worker	27	3	27	20	26	21
Medical staff	5	7	11	12	0	7
Other	13	1	2	1	2	5

4.15. Trends in condom use among service women selling sex in Vientiane

Vientiane is the only city that always included in all surveillance round. To ensure comparability across survey the trends were constructed exclusively for Vientiane. The trend in condom use at last sex with clients has remained around 90% since the first round of surveillance in 2001. Consistent use of condoms with client in the past 3 months remained unchanged 2008 to 2009.

Figure 3: Trend in condom use with clients among service women in Vientiane



V. Conclusions and recommendations

5.1. Conclusions

The level of risk behaviors among service women has remained unchanged since 2008. There is no significant difference in behaviors between service women from remote areas and those from Vientiane capital. However, access to services may vary significantly by province. Uptake of VCT services and STI treatment remain too low in some province.

5.2. Recommendations

- Due to the frequent mobility of service women and their short duration staying in one establishment, prevention interventions should be done regularly.
- Prevention should include condom skills to address the frequency of condom breakage and inadequate use of condoms
- Prevention should include condom negotiation skills with focus on new

- Messages on condom use with regular clients should be strengthened.
- Distribution of condoms to service women is insufficient to bolster the importance messages on consistent use of condoms with clients. The strategy on condom distribution needs to be revised.
- Female condoms appear acceptable and the use of female condom as an alternative for client refusing to use condoms should be piloted.
- Geographical coverage should be strategic to ensure quality services in all provinces.
- Quality of implementation should be strengthened in provinces to ensure quality services.
- Each province should have a STI service dedicated to FSW (private, NGO, DIC....). Drop-in centers appear widely used where they exist and they should therefore be sustained.
- Reproductive health services (family planning) should be integrated with STI services to address the issue of unwanted pregnancies and frequent abortions among service women.
- Regular partners of service women (i.e. regular clients) need to be treated for STI.
- Results of the present survey should be shared with the Lao Women Union to address issues related to violence.
- Gaps in surveillance:
 - Ethnographic study among FSW on mobility and motivations.
 - Inclusion of HIV module questions in DHS.
 - Assessment of use and satisfaction of female condom.



Table

Table 1: Sentinel sites for service women by year of survey

Table 2: Demographic characteristics of service women, by province.....

Table 3: Age at first sex and first sexual partner among service women.....

Table 4: Reproductive history, use of contraceptives and history of abortion among service women

Table 5: History of selling sex and mobility of service women

Table 7: Characteristics of last client

Table 8: Condom availability

Table 9: Condom use.....

Table 10: Use of female condoms.....

Table 11: Report of aggressions

Table 12: Type of drug use among Service Women.....

Table 13: Knowledge of HIV transmission and prevention among service women

Table 14: Reported symptoms of STI and attendance to routine checkups

Table 15: Uptake of HIV testing services

Table 16: Array of services received by service women at last HIV testing

Table 17: Coverage of prevention services

Figure

Figure 1: Condom breakage past 3 months by use of two condoms at a time

Figure 2: Source of treatment at last episode of STI, which occurred in the past three months

Figure 3: Trend in condom use with clients among service women in Vientiane.....



Content

I. Background and Rationales.....	1
II. Objectives.....	2
2.1. Primary objective:	2
2.2. Secondary objectives:	2
III. Methods	2
3.1. Survey design:	2
3.2. 3.2. Survey population:	2
3.3. Survey areas:	3
3.4. Sample sizes:	3
3.5. Sampling design:	3
3.6. Survey procedures:	4
3.7. Ethical review:	4
IV. Results	5
4.1. Demographic characteristics	5
4.2. First sex	6
4.3. Reproductive history, contraceptive use and abortions	7
4.4. History of selling sex	8
4.5. Sexual Partners	9
4.6. Last clients characteristics	10
4.7. Availability and use of male latex condoms	10
4.8. Female condom	13
4.9. Violence to service women	14
4.10. Drug use	14
4.11. Knowledge of HIV	15
4.12. Reported symptoms of STI and attendance to STI clinic	16
4.13. Uptake of HIV testing services	18
4.14. Source of information and coverage of prevention services	19
4.15. Trends in condom use among service women selling sex in Vientiane	20
V. Conclusions and recommendations	21
5.1. Conclusions.....	21
5.2. Recommendations	21
Reference.....	23



Survey Team for BSS among Service Women in 5 provinces

From CHAS:

Dr. Chansy Phimpachanh	Director of CHAS
Dr. Phouthone Southalack	Deputy Director of CHAS
Dr. Chanthone Khamsibounheuang	Deputy Director of CHAS
Dr. Keophouvanh Duangphachanh	Chief of Administrative Unit
Dr. Khanthanouvieng Sayabounthavong	Chief of STI Unit
Dr. Ratthiphone Oula	Deputy Chief of STI Unit
Dr. Beuang Vangvanh	Chief of Planning Unit
Dr. Phengphet Phetvixay	Chief of IEC Unit
Dr. Panina Phoumsavanh	Deputy Chief of Adm Unit
Dr. Ketmala Banchongphanit Unit	Acting Chief of Surveillance
Dr. Phutthaly Keomukda	Deputy Chief of IEC Unit
Dr. Khanti Thongkham	Technical staff from Surveillance Unit
Dr. Buathong	Technical staff from Planning Unit
Dr. Souphanthong Manichanh	Technical Staff from STI Unit
Dr. Vanvilay Sphabmixay	Technical Staff from Surveillance Unit
Dr. Chanthasouk Bansalith	Technical Staff from STI Unit
Dr. Chanhom Phinsavanh	Technical Staff from IEC Unit
Dr. Chanvilay Thammachak	Technical Staff from surveillance Unit
Ms. Southaphone chittaphong	Technical Staff from Surveillance Unit

From Vientiane Capital:

Dr. Inlavanh Keobounphanh Dept	Director of Vientiane Capital Health
Dr. Phonepaseut Ounaphom	PCCA sec of VTC
Dr. Hapheng Phommalath	Head of Sikhottabong District Health Office
Dr. Khamchanh Sayavong	Head of Chanthabouly District Health Office
Dr. Khanthalat Philaphandeth Office	Head of Xaysettha District Health
Dr. Bounthanong songchaksinh Office	Head of Xaythani District Health
Dr. Lavanh Vongsavanthong	PCCA sec of VTC
Dr. Xayavong Duanmany	PCCA sec of VTC
Dr. khankeo Somvang	DCCA Xaythani, VTC
Ms. Viengxay Chanthavong	DCCA Xaythani, VTC
Dr. Saysavath Chanthavong	DCCA Xaythani, VTC
Ms. Vilayphone Chanthakhoun	DCCA Xaythani, VTC
Ms. Souphalack Bounsom	DCCA xaythani, VTC
Mr. Bounheuang Manivong	DCCA Chanthabouly, VTC
Dr. Vilavanh Viengxayavong	DCCA Chanthabouly, VTC
Dr. Phitsamay Manivong	DCCA Chanthabouly, VTC
Ms. Malayphone simmavong	DCCA Chanthabouly, VTC

From Saravane:

Dr. Oudomsack Sisana
Ms. Chanthaly sisoulat
Dr. Yommala
Ms. Sengthong Keovongkot

PCCA sec of Saravane
PCCA sec
Clinician from Saravane Prov. Hosp
PCCA sec

From Attapeu:

Dr. Phetlatda Phoumsavanh
Dr. Viengsay
Ms. Keoduangchit
Ms. Bouasavanh Phothisane

PCCA sec
PCCA sec
PCCA sec
Education sector

From Luang namtha:

Dr. Khamchanh Inthongxay
Mr. Somlit Senevanhpenh
Dr. Keo Inthavong
Dr. Sengvong Bouttavong
Ms. Dengthong Losomto
Ms. Thoumy Vongbouasy
Ms. Monekeo Chanthavong
Ms. Pinhkeo Boutthavong

Deputy Director of Health Department
Deputy Director of Health Department
Adm unit
PCCA sec
PCCA sec
PCCA sec
PCCA sec
PCCA sec

From Bokeo:


Dr. Khamla
Ms. Khammanh
Dr. Phouthone
Ms. Bouathip
Ms. Aksone

PCCA sec
PCCA sec
PCCA sec
PCCA sec
PCCA sec





ACRONYMS AND ABBREVIATIONS	
AIDS	Acquired Immunodeficiency Syndrome
BSS	Behavioral Surveillance Survey
CLE	Center for Laboratory and Epidemiology
FSW	Female sex worker
HIV	Human Immunodeficiency Virus
NCCA	National Committee for the Control of AIDS
NGO	Non-governmental organization
PCCA	Provincial Committee for the Control of AIDS
PDA	Personal digital assistant PDR People's Democratic
Republic	
STI	Sexually transmitted infection
VCT	Voluntary Counseling and Testing



I. Background and Rationales

Lao PDR has conducted three rounds of behavioral and biological surveillance among Service Women at sentinel sites in 2001, 2004 and 2008. Provinces included in the sentinel surveillance were the most populated and those with easy access to the target populations. The surveillance schedule was adapted in 2009, in order to conduct regular surveys in a small number of provinces to serve as early warning and to closely monitor progress of programs implemented to this target population .

Women exchanging sex for money take diverse forms in Lao PDR. They are indirect sex workers as Service Women working in hotels, guest houses, night clubs, karaoke, beer bars, and small drink shops . There is an increasing reporting of women contacting their clients through phone calls . Indeed, 71% of the male clients surveyed in 2009 had met their paid sexual partners at drink shops or night-clubs. Other places where clients met FSW were guest houses and hotels, which were named by 15% of the clients in 2008.

Service women are female sex workers characterized by a low number of clients, a short duration of selling sex and a high level of condom use . Condom use with both clients and non-commercial non-regular partners has dramatically increased since 2004. These specificities largely explain why the prevalence of HIV has not increased in this group of female sex workers and may even be decreasing. Fortunately, very few service women use drug and when they do, they do not inject. The short duration of working at a specific place as well as the short duration of selling sex requires intensified prevention which is highlighted by the insufficient knowledge of HIV transmission reported by the majority of service women. Although condoms appear accessible, few women purchase their own condoms and most get their condoms from prevention projects or drink shop managers.

Service women are young and young women are biologically more vulnerable to STI than their peers. STI prevalence serves as a biological proxy for unprotected sex. The prevalence of STI among service women has dramatically decreased in provinces where periodic presumptive treatment (PPT) was implemented, which provinces also achieved a high level of condom use in commercial sex.

The integrated biological and behavioral survey conducted in 2008 as part of Lao PDR national surveillance found a 0.4% prevalence HIV among service women and 0.3% among state enterprise electricity workers (half of whom had purchased sex from service women in the year prior to survey). While syphilis prevalence was also low (0.4% life-time), the prevalence of other curable STI remained high among service women (Chlamydia 18% and gonorrhea 7%). Chlamydial infections are also prevalent even in the general population (10%).
Reducing



rates of curable STI is a priority for HIV prevention strategy to countries with low-level HIV epidemics, where targeted interventions for vulnerable populations appear to be efficient and cost effective.

A set of indicators detailed in the grant assistance paper is required to document the end of the five years project of Global Fund Round 1, coming to a term in 2009. These indicators includes: trend in HIV prevalence, knowledge of HIV prevention, condom use, and uptake of STI services.

II. Objectives

2.1. Primary objective:

- To determine the level of HIV-related risk behaviors among service women

2.2. Secondary objectives:

- To provide data to construct trends in prevalence of sexual behaviors and uptake of prevention services
- To estimate the coverage and outcomes of the national HIV and STI prevention strategy
- To determine health-seeking behaviors among those reporting symptoms of STIs currently or/and in the previous year



III. Methods

3.1. Survey design:

A cross-sectional study design was used. Participation to the survey was voluntary and anonymous.

3.2. Survey population:

Defining and identifying women who sell sex for money in Lao PDR can be particularly difficult. Women working in small drink shops and nightclubs may engage in commercial sex transactions, but their employment in these venues does not automatically signify that they are selling sex, as opposed to just serving beer or having conversation with their customers. This population has been used in previous round of surveys to approximate female sex workers. However, women were not screened to determine whether they sold sex or not. Instead, all women who worked in these establishments and had direct contact with clients, whether by selling them drinks or sitting with them, were defined as service women.

Those eligible were:

- Women
- Aged between 15 to 49 years old
- Working in small drink shops, beer gardens, karaoke bars, or other entertainment establishments
- Not hanged over at the time of the interview

3.3. Survey areas:

Service women were selected in the capital city of selected provinces. The provinces included in the 2009 survey were selected purposively. The provinces selected included some provinces covered by the previous surveillance rounds to allow for comparability and construction of trends over time, as well as provinces that had never been surveyed in the past to explore potential variations in behaviors. Provinces surveyed in 2009 were: Vientiane capital, Bokeo, Luang Namtha, **Saravane**, and Attapeu.

The provinces included in previous and present round of second generation surveillance by target population are as follows:

Table 1: Sentinel sites for service women by year of survey

AREA	2001	2004	2008	2009
Vientiane municipality	○	○	○	○
Champassak		○	○	
Savannakhet	○	○	○	
Bokeo		○	○	○
Luang Prabang	○		○	
Luang Namtha		○	○	○
Saravane				○
Attapeu				○

3.4. Sample sizes:

Sample size was set at 300 for Vientiane capital and 150 for each of the remaining province.

3.5. Sampling design:

The Provincial Committee for the Control of AIDS Secretariat (PCCAsec) of the surveyed provinces established the sampling frames by listing all identified small drink shops, night clubs, guest house and karaoke bars located in the survey area with an estimate of the number of women working at each location. Establishments were selected randomly proportional to size. At second stage women were selected randomly in order to include:

- 20 participants from selected night clubs
- 5 participants from selected restaurants / drink shop or karaoke bars
- All participants when the provincial sample size was inferior to the number of women enumerated (**Saravane** and Bokeo).