

FAMILY HEALTH RESEARCH

A forum for putting knowledge into practice

INSIDE

- 2 | Clinical research update
- 4 | Evaluating tenofovir gel in South Africa
- 6 | Participating in microbicide trials
- 7 | Preparing for microbicides
- 8 | Research and advocacy partnerships



MICROBICIDES FOR HIV PREVENTION

This issue describes efforts to develop a safe, effective, and acceptable microbicide to prevent HIV.

Scientists around the world are working diligently to develop new ways to prevent HIV, including vaccines, pre-exposure prophylaxis (when drugs normally used to treat HIV are used for prevention), and topical microbicides. Advocacy and support for prevention research are strong, and the availability of an effective product—although likely years away—could help alleviate the HIV epidemic in sub-Saharan Africa and worldwide.

The Joint United Nations Programme on HIV/AIDS recently announced that the prevalence of HIV has stabilized or declined in many parts of sub-Saharan Africa. Nevertheless, sub-Saharan Africa still suffers the most from the epidemic. More than two-thirds of all HIV-

infected people in the world live in this region. Of these, more than 60 percent are women. Young women are even more disproportionately affected. In South Africa, for example, women make up 90 percent of the HIV-infected population of 15- to 24-year-olds.

Women are at especially high risk of HIV infection because of biologic susceptibility, economic dependence on male partners, and inability to negotiate condom use within all of their sexual relationships. A method of HIV prevention that a woman could control, such as a topical microbicide, could help overcome some of these inequities and save many lives.

Both vaginal microbicides (for use by women) and rectal microbicides (for use by women and men) are under development. This issue will focus on the vaginal products that have reached clinical trials of effectiveness in humans. An effective product could be a powerful addition to current efforts to protect against HIV, including abstinence, mutual monogamy with uninfected partners, condom use, treatment of sexually transmitted infections, and male circumcision.

In July 2011, FHI became FHI 360.



FHI 360 is a nonprofit human development organization dedicated to improving lives in lasting ways by advancing integrated, locally driven solutions. Our staff includes experts in health, education, nutrition, environment, economic development, civil society, gender, youth, research and technology – creating a unique mix of capabilities to address today's interrelated development challenges. FHI 360 serves more than 60 countries, all 50 U.S. states and all U.S. territories.

Visit us at www.fhi360.org.

CLINICAL RESEARCH UPDATE

KEYPOINTS

- An effective vaginal microbicide has not yet been identified.
- Three products are being studied in large-scale effectiveness trials.
- The newest candidates are designed to stop HIV from replicating.

The research pipeline is filled with hopeful microbicide candidates.

Research to develop an effective vaginal microbicide began nearly three decades ago. Today, more than 60 different products are in some stage of laboratory or clinical development. Eleven of these products are in human clinical trials, and three are in large-scale effectiveness trials. However, no candidate microbicide has yet been shown to prevent HIV or other sexually transmitted infections (STIs) in humans.

“The process of product development is long, but the HIV-prevention community is hopeful that an effective topical agent will be identified within the next decade,” says Dr. Willard Cates, president of research at FHI.

Types of products

The “first generation” of microbicide candidates are typically formulated as gels, which are applied to the vagina before each act of sexual intercourse. These candidates employ one of three mechanisms that should theoretically prevent HIV and other STI pathogens from infecting cells. Some are surfactants (agents that reduce the surface tension of a liquid) that inactivate a bacterium or virus by damaging its surface membrane. Others enhance the vagina’s natural defenses against pathogens, for example by maintaining the natural acidity of the vagina in the presence of semen, which is alkaline. Still others block viruses from attaching to healthy cells.

The next generation of microbicide candidates is based on antiretroviral drugs. These products specifically target HIV, acting later in the viral life cycle to prevent HIV from replicating once it has entered the body. Although some are formulated as gels, others may be released through vaginal rings. And, unlike the first-generation candidates, some of the newer products may be delivered on a daily or monthly basis (such as through a vaginal ring) regardless of the timing of intercourse.

Eventually, a combination of different products with more than one mechanism of action may be the most effective way to prevent a wide range of STIs. Products with a contraceptive ability will also be important for meeting diverse reproductive health needs.

Research process

After a microbicide candidate is identified, its safety and effectiveness is first assessed in test tubes and animal models. If the product passes this stage of testing, it enters human clinical trials.

In Phase I microbicide trials, scientists assess the safety of the product, determine its acceptability among potential users, and identify its appropriate dose and formulation. In Phase II trials, scientists again assess safety and acceptability, but among more participants and for a longer period.

The final stages of microbicide trials (Phase IIb and Phase III trials) are used to evaluate a product’s effectiveness in preventing HIV and other STIs. Phase IIb trials are typically smaller than Phase III trials, giving some indication of effectiveness and helping scientists decide whether a larger Phase III trial is necessary. Lasting up to four years, Phase III trials enroll thousands of participants who are at risk of HIV, such as those living in communities with a relatively high incidence of HIV.

Although all participants in microbicide trials are regularly counseled on how to reduce their risks of acquiring HIV and other STIs, prior experience has shown that even with optimal counseling and free access to condoms, most women are not able to negotiate condom use every time they have sex. The effectiveness of a candidate microbicide is tested in those sex acts in which women are not able to use condoms.

Ruling out candidates

Three first-generation products—SAVVY, cellulose sulfate, and Carraguard—were ruled out as effective microbicides in advanced clinical trials.

In 2005, FHI closed its Phase III trial of SAVVY (a surfactant) in Ghana when an independent data monitoring committee determined that



A counselor explains condom use to a woman participating in a microbicide trial.

the incidence of HIV was so low that the study would not be able to determine whether SAVVY could prevent HIV. FHI closed its parallel Phase III trial of SAVVY in Nigeria in 2006, after a similar committee concluded that the trial was unlikely to find a protective effect if it continued. Earlier studies had shown that spermicides containing another surfactant, nonoxynol-9, may actually increase the risk of HIV infection when women at high risk of infection use them frequently.

In 2007, CONRAD closed its Phase III trial of cellulose sulfate (an inhibitor of viral attachment and entry) in South Africa, Benin, Uganda, and India after a planned interim analysis suggested that the product might contribute to an increased risk of HIV. Because of these safety concerns, FHI simultaneously closed its Phase III trial of cellulose sulfate in Nigeria. However, the final data from both trials showed that cellulose sulfate had not significantly affected the risk of HIV infection either positively or negatively.

Most recently, the Population Council completed a Phase III trial of Carraguard (another inhibitor of viral attachment and entry) in

South Africa. Results, released in 2008, showed that the Carraguard gel was safe and acceptable. However, it did not reduce the risk of acquiring HIV. Although these results were disappointing, the Carraguard trial was the first effectiveness trial to be completed as planned for a product that was developed specifically as a microbicide—an historic accomplishment in the eyes of many microbicide advocates.

More effectiveness trials

Three microbicide candidates—BufferGel, PRO 2000, and tenofovir—are currently in large-scale effectiveness trials. Three additional products—ACIDFORM, the invisible condom, and dapivirine—will enter the advanced stages of clinical trials soon (see table).

“As with all other HIV-prevention approaches, no vaginal product is expected to offer 100 percent protection against HIV or other STIs,” says Dr. Cates. “At best, an effective product would prevent infection perhaps half the time. But even a partially effective microbicide could help women who might otherwise not be able to protect themselves at all.”

MICROBICIDE EFFECTIVENESS TRIALS

Mechanism of action	Microbicide	Trial status
Enhances vaginal defenses against pathogens	BufferGel*	The Microbicide Trials Network is conducting a Phase II/IIb trial of BufferGel vs. 0.5% PRO 2000 gel vs. a placebo vs. no intervention (condoms only) in Malawi, South Africa, Zambia, Zimbabwe, and the United States. Results of the trial, begun under the HIV Prevention Trials Network, are expected in 2009.
	ACIDFORM*	CONRAD and partners are planning a Phase III trial of a diaphragm with ACIDFORM gel in Madagascar (for the prevention of STIs other than HIV).
Inhibits viral attachment and entry	PRO 2000*	The Microbicides Development Programme is conducting a Phase III trial of 0.5% PRO 2000 gel vs. a placebo in Zambia, Tanzania, and South Africa. Results are expected in 2009. See “BufferGel” above for information on a second trial involving PRO 2000.
	Invisible condom*	Scientists at Laval University and partners are planning a Phase III trial of the invisible condom (a gel containing sodium lauryl sulfate, a common ingredient in soap). Sites are to be determined.
Inhibits viral replication	Tenofovir	The Centre for the AIDS Programme of Research in South Africa (CAPRISA) and partners are conducting a Phase IIb trial of 1 percent tenofovir gel vs. a placebo in South Africa. Results are expected in 2010. The Microbicide Trials Network is planning a Phase IIb study of tenofovir gel vs. oral tenofovir vs. oral Truvada (a combination of tenofovir and a second antiretroviral drug, emtricitabine) vs. a placebo gel vs. an oral placebo in South Africa, Malawi, Uganda, Zambia, and Zimbabwe.
	Dapivirine	The International Partnership for Microbicides is planning a Phase III trial of dapivirine gel. Sites are to be determined.

* Also being evaluated for contraceptive ability.

EVALUATING TENOFVIR GEL IN SOUTH AFRICA

Community members play important role in trial.

The Centre for the AIDS Programme of Research in South Africa (CAPRISA) and its partners are testing the effectiveness of tenofovir gel—a promising antiretroviral-based microbicide—in two South African communities. The intense collaboration between CAPRISA and the community of Vulindlela demonstrates just how vital community involvement can be in the preparation and conduct of a microbicide trial.

“Vulindlela is one of many communities we have worked in,” says Dr. Quarraisha Abdool Karim, CAPRISA’s associate scientific director and one of the principal investigators of the tenofovir gel trial. “But our experience in Vulindlela has been unique.”



The semi-rural district of Vulindlela, shown here, is one of two communities participating in the trial.

A semi-rural district outside of Durban, Vulindlela is home to almost half a million people. For years, the residents of Vulindlela had experienced the effects of AIDS on their community, watching as family graveyards filled up at an astonishing pace. Finally, the traditional village chief, or *Inkosi*, was compelled to take an unusual step: He contacted a group of local scientists.

Dr. Abdool Karim was one of the scientists who met with the traditional leaders of Vulindlela one Saturday morning in 2001. After welcoming the visitors, the *Inkosi* apologized that he could not spend the entire day with the group, as planned. “He had several funerals to attend,” recalls Dr. Abdool Karim.

In response to the concerns of the traditional leaders, CAPRISA conducted extensive research on the nature of the HIV epidemic in the community. For two years, scientists hosted focus group discussions with local stakeholders, studied the HIV cases from local clinics, and conducted “verbal” autopsies to establish the main causes of death in the community. Armed with these data,

THE TRIAL . . .

- Phase IIb study evaluating the safety and effectiveness of tenofovir gel in preventing HIV infection
- Enrolling 980 sexually active, HIV-uninfected women, ages 18 to 40, who are at high risk of acquiring HIV
- Being conducted at clinical research sites in Vulindlela and Durban
- Study participants randomized to use either 1 percent tenofovir gel or a placebo gel
- Participants counseled to apply the gel via single-use, prefilled applicators before and after intercourse
- Participants given free condoms and counseled on reducing the risks of infection
- Participants and their partners provided with free screening and treatment for sexually transmitted infections
- Results expected in 2010

. . . AND WHY IT IS UNIQUE

- First effectiveness trial of an antiretroviral-based microbicide
- First microbicide trial that an institution in the developing world is leading
- First microbicide trial that a South African government agency is co-funding
- First time that South Africa has secured an up-front voluntary license to manufacture and distribute a microbicide if it is proven effective



they were able to secure funding to establish an AIDS treatment program. Today, the program cares for more than 5,000 patients, including 1,300 who are receiving antiretroviral treatment. As part of a community fellowship program, CAPRISA also trained 32 community members as HIV/AIDS educators.

Despite these accomplishments, the prevalence of HIV remained high in Vulindlela. At last measure, about 55 percent of women ages 20 to 24 were infected. This alarming statistic highlighted the importance of finding an effective HIV-prevention method that women could control. Ultimately, CAPRISA selected Vulindlela as one of the study sites for its tenofovir gel trial, which began in May 2007.

Together, the researchers and local traditional leaders established the Vulindlela Community Research Support Group to function as a liaison between the

community, study participants, and research team. The group is a mixture of community leaders, representatives from religious and women's organizations, young people, government health care providers, and other stakeholders who review study materials and provide advice on recruitment and retention. They also function as a conduit through which community members can voice any concerns about the trial or other research that CAPRISA is conducting.

So far, more than 3,000 women in Vulindlela have received general information about the tenofovir gel trial. Among these women, 800 volunteers have been screened, and 454 have been enrolled in the study. As a testament to CAPRISA's dedication to building research capacity in the region, all of the study staff—including the recruitment teams, community educators, and project managers—are South African.

CAPRISA

CAPRISA is a nonprofit organization dedicated to researching new ways to prevent and treat HIV and tuberculosis, and to building local research infrastructure and capacity. CAPRISA is collaborating with CONRAD, FHI, and the South African Ministry of Science and Technology (through LIFE/lab) on its effectiveness trial of tenofovir gel. More information about the trial is available at: <http://www.caprisa.org/Projects/microbicides.html#5>.

ADHERENCE

A participant's adherence to the study regimen is absolutely vital for valid results in a trial. Yet, unstable living conditions and other stressors sometimes make it difficult for women to follow the directions for using the study products.

The CAPRISA team has implemented an innovative Adherence Support Program to help increase adherence rates. The program is based on the intention-motivation-behavioral skills model of HIV prevention, which reduces the risk of infection through behavioral change.

Before enrolling in the trial, potential participants learn about using the gel applicators, possible side effects of the gel, the time required to participate in the trial, and the benefits and risks of participating. Once enrolled, each woman receives tailored counseling provided by a nurse using an approach called "motivational interviewing." Counselors use various teaching techniques to emphasize that the gel has to be inserted

within 12 hours before sex and again up to 12 hours after sex, no matter how many times a woman has sex within that period. This core message is summarized as BAT-24: a dose **B**efore sex, a dose **A**fter sex, and not more than **T**wo applications within **24** hours.

When a woman reports problems with adherence, the nurse helps her develop a plan to address them. The counselor also writes "adherence support prescriptions" that spell out practical, customized suggestions for each individual's situation.



Explicit instructions on using the gel applicators, combined with tailored counseling during the trial, help participants adhere to the study regimen.

PARTICIPATING IN MICROBICIDE TRIALS

KEYPOINTS

- The informed consent process aims to give participants a complete, continuing understanding of a trial.
- Community representatives can help researchers design trials that are acceptable, relevant, and ethical.
- Ensuring continuity of care for women who become infected with HIV during a trial is challenging.



Community nurses, shown here with FHI's Stella Kirkendale, are working closely with microbicide trial participants in Malili, Malawi, to address any concerns the women have about the trial.

Researchers use multiple strategies to protect participants.

Since 1996, more than 20,000 women have participated in clinical trials to help identify a safe and effective vaginal microbicide.

All of the women volunteered for the trials, knowing they were going to help answer a question about the safety, acceptability, and in some cases effectiveness of a study product. They agreed to return to their study sites every three months, or even monthly, to receive counseling, supplies of gel and condoms, and an HIV test.

Study staff also made a commitment to the women. In their study protocols, counseling messages, and educational materials, the staff outlined how they would protect the safety, confidentiality, and human rights of each study participant. Their experiences have shown just how complex—and critical—that commitment can be.

Truly informing women

First, researchers have an ethical responsibility to ensure that participation in a clinical trial is voluntary and that the women who enroll in a trial understand exactly what their participation means.

The women must understand that the researchers do not know whether the study product works, that a portion of the participants may be randomly assigned to use a placebo that contains no microbicide, and that all participants need to continue using condoms and other HIV-prevention measures throughout the study.

Explaining technical terms and research concepts such as randomization and placebos to potential participants can be difficult. As a result, researchers have developed creative materials such as illustrated booklets, videos, and interactive computer-based educational programs to help explain them. Many trials

also use a questionnaire to assess a woman's understanding of a study before she signs an informed consent form agreeing to participate in the study.

However, researchers also recognize that informed consent should be a continuing process rather than a one-time event. In the two Phase III cellulose sulfate trials that have been conducted, for instance, the study staff administered this type of questionnaire both before and during the trials.

Involving communities

Researchers also have an increasing appreciation for the value of involving communities in decisions about how the trials are designed and implemented. Most trials have community liaison officers, and many have community advisory boards that serve as a link between researchers and the communities where the research is conducted. Some trials also train participants to educate their fellow participants and other community members about the study.

These community representatives help ensure that the trials are designed and implemented in an acceptable, relevant, and ethical way. The representatives review study protocols and informed consent forms, evaluate draft educational materials, relay community concerns to the study staff, refute rumors, and help disseminate study results. Such community involvement generates support for microbicide research and reduces the likelihood that a woman will be stigmatized because of her participation in a trial.

Ensuring access to care

One of the most complex issues facing microbicide researchers is how to provide access to treatment for the relatively small number of women who seroconvert (become infected with HIV) during a trial, as well as for the many more volunteers who are ineligible to participate because pretrial screening showed that they were already infected.

Funding for microbicide trials does not always include support for antiretroviral therapy (ART) or other services for people living with HIV. However, cost is not the only barrier to treatment. The complexity of ensuring

continuity of care over a person's lifetime is also a challenge for research organizations conducting short-term studies.

A 2006 study by the Global Campaign for Microbicides found that the standard of HIV care for participants who seroconvert during microbicide trials has varied. One trial set aside funds to support five years of ART for every woman who seroconverted during the trial. Others offered to enroll seroconverters into trials of ART regimens. Most referred the women to the closest source of HIV care, and many were able to establish written agreements with programs that provide free ART.¹ A study by FHI suggests that such referrals are acceptable to study participants.²

The Global Campaign for Microbicides recommends that microbicide trials be conducted at or in partnership with local HIV care centers, and that researchers use their trials as opportunities to strengthen local standards of care for people living with HIV.

References

- 1 Heise H, Philpott S, Shapiro K, et al. Mapping access to HIV care in microbicide effectiveness trials. *Microbicides* 2008, New Delhi, India, February 24–27, 2008.
- 2 MacQueen KM, Namey E, Chilongozi DA, et al. Community perspectives on care options for HIV prevention trial participants. *AIDS Care* 2007;19(4):554–60.

PROGRAM DISPATCH

PREPARING FOR MICROBICIDES

Introducing new products is complex.

As clinical trials continue, scientists and the broader microbicide community are also preparing for the complex process of introducing a vaginal product once it has been shown to be effective.

Foremost, scientists are evaluating whether women and men will actually use a microbicide if it becomes available. Qualitative research on a product's acceptability may help to inform the design and development of new products, estimate how well women will adhere to an eventual microbicide, and influence how microbicides might one day be marketed.

Most acceptability studies focus on product characteristics such as formulation, color, texture, and the type of application. Others have investigated the importance of contraceptive ability among potential consumers, the influence of sexual partners on product use, and other factors that could differ between individuals or communities.

Few studies have evaluated the role that health care providers would play in promoting microbicides.¹ Yet, if and when a vaginal

microbicide becomes available, health professionals would strongly influence whether women could access the product and how they would use it.

To help ensure the acceptability and use of microbicides outside of the research setting, the microbicide community is also:

- Considering the cost and affordability of products for potential consumers, reproductive health programs, and donors
- Identifying the most appropriate targets for initial introduction
- Developing counseling messages and deciding how microbicides would fit into a larger HIV-prevention strategy
- Determining appropriate service-delivery points and evaluating the capacity of service-delivery systems and health care providers to offer the products
- Collaborating with donors, regulatory bodies, and other stakeholders who could influence introduction, since most of the products would be manufactured and distributed in developing countries without the help of large pharmaceutical companies

Reference

- 1 Mantell JE, Myer L, Carballo-Diéguez A, et al. Microbicide acceptability research: current approaches and future directions. *Soc Sci Med* 2005;60(2):319–30.

Resources

Informed Consent in HIV Prevention Trials: Report of an International Workshop.
www.popcouncil.org/pdfs/ICWorkshop.pdf

Partnering for Care in HIV/AIDS Clinical Trials: A How-To Manual.
Order a copy of this soon-to-be-released publication by writing to publications@fhi.org.

WHO NEEDS TO BE READY

- Consumers
- Health professionals
- Reproductive health programs
- Service-delivery systems

FAMILY HEALTH RESEARCH is a newsletter of the FHI Contraceptive and Reproductive Health Technologies Research and Utilization (CRTU) program, which is supported by the U.S. Agency for International Development (USAID) under the terms of Cooperative Agreement No. GPO-A-00-05-00022-00. The contents do not necessarily reflect the views of USAID.

Managing Editor:
Kerry Wright Aradhya

Writers:
Kathleen Henry Shears, Jill Moffett

Art and Production Editor:
Karen Dickerson

Photos:
Cover: Karen Tweedy-Holmes/
The Population Council, Inc.

Page 2: Bev Symmonds/
The Population Council, Inc.

Page 4: CAPRISA;
Elizabeth T. Robinson/FHI

Page 5: Elizabeth T. Robinson/FHI

Page 6: Michael Szpir/FHI

Page 8: Microbicide Trials Network

Inclusion of persons in photos
should not be construed as
indicating their HIV status.



USAID
FROM THE AMERICAN PEOPLE



FHI Headquarters
PO Box 13950
Research Triangle Park, NC 27709
USA
Tel: 1.919.544.7040
www.fhi.org

Send comments to:
[Kerry Wright Aradhya](mailto:Kerry.Wright.Aradhya@fhi.org)
familyhealthresearch@fhi.org

July 2008
Volume 2, Issue 2

RESOURCES

RESEARCH AND ADVOCACY PARTNERSHIPS

Microbicide Trials Network

This clinical trials network is dedicated to developing and evaluating an effective microbicide. Funded by the U.S. National Institutes of Health, the network includes a core center of operations (supported by the University of Pittsburgh, FHI, and others) and 18 clinical research sites in seven countries. Most recently, the network conducted a Phase II trial in India and the United States showing that tenofovir gel is safe and acceptable for daily use. More information about the network and its trials is available at: <http://www.mtnstopshiv.org/>.

Microbicides Development Programme

Based in the United Kingdom, this partnership of organizations in Europe and Africa primarily conducts research to develop an effective vaginal microbicide. Its research includes studies to determine mechanisms of action for different products, preclinical evaluations of potential microbicides, and clinical trials of promising candidates. The partnership is currently conducting a Phase III trial of PRO 2000 gel in Africa. Learn more at: <http://www.mdp.mrc.ac.uk/>.

International Partnership for Microbicides

This global collaboration between public- and private-sector organizations promotes

the rapid development and availability of microbicides. The partnership also conducts its own research, focused on developing and testing antiretroviral-based microbicides such as dapivirine gel. Learn about the partnership's clinical trials and how it supports access to future microbicides at: <http://www.ipm-microbicides.org/>.

Alliance for Microbicide Development

This nonprofit organization is composed of representatives of more than 200 biopharmaceutical companies, research organizations, and advocacy groups dedicated to accelerating the development of a safe, effective, and affordable microbicide. One of its many activities is to monitor and analyze the current pipeline of microbicide candidates. View more about the alliance, including an up-to-date database of information on microbicide research and development, at: <http://www.microbicide.org/>.

Global Campaign for Microbicides

This network of more than 285 nongovernmental organizations uses international advocacy, policy analysis, and social science research to build support for microbicides and other HIV-prevention options for women. The campaign's regional activities include raising awareness of microbicides and protecting the needs and interests of trial participants and other potential users. Find out about the campaign's work at: <http://www.global-campaign.org>.



The Microbicide Trials Network—whose clinical research staff in Zambia are shown here—is one of several partnerships dedicated to developing and building support for microbicides.